

Introduction to Transient Analysis

Prof. John C.S. Lui
Department of Computer Science & Engineering
The Chinese University of Hong Kong

Outline

- 1 Motivation
- 2 General Solution
- 3 Uniformization Method
 - Uniformization
 - Probabilistic Interpretation
- 4 Practical Issues
 - Finite Sum
 - Extension to non-homogeneous CTMC

Outline

- 1 Motivation
- 2 General Solution
- 3 Uniformization Method
 - Uniformization
 - Probabilistic Interpretation
- 4 Practical Issues
 - Finite Sum
 - Extension to non-homogeneous CTMC

Motivation

- Motivation of why we need to perform *transient analysis*.
- Important questions are:
 - state of the model at the end of a time interval,
 - the time until an event occurs,
 - the residence time in a subset of states during a given interval,
 - the number of given events in an interval.

Outline

- 1 Motivation
- 2 General Solution**
- 3 Uniformization Method
 - Uniformization
 - Probabilistic Interpretation
- 4 Practical Issues
 - Finite Sum
 - Extension to non-homogeneous CTMC

Let $\{X(t), t \geq 0\}$ be a CTMC with finite state space $\mathcal{S} = \{s_i : i = 1, \dots, M\}$ and \mathbf{Q} the transition rate matrix:

$$\mathbf{Q} = \begin{bmatrix} -q_1 & q_{12} & \cdots & q_{1M} \\ q_{21} & -q_2 & \cdots & q_{2M} \\ \vdots & \vdots & \cdots & \vdots \\ q_{M1} & q_{M2} & \cdots & -q_M \end{bmatrix} \quad (1)$$

where $q_i = \sum_{j=1, j \neq i}^M q_{ij}$.

Let $\mathbf{\Pi}(t)$ be a $M \times M$ matrix where $\pi_{ij}(t)$ in $\mathbf{\Pi}(t)$ is :

$$\pi_{ij}(t) = P[X(t) = s_j | X(0) = s_i] \quad (2)$$

Based on Kolmogorov's forward equation, we have:

$$\mathbf{\Pi}'(t) = \mathbf{\Pi}(t)\mathbf{Q} \quad (3)$$

Solving the above matrix equation, we have:

$$\mathbf{\Pi}(t) = \mathbf{e}^{\mathbf{Q}t} \quad (4)$$

Let $\pi(t) = [\pi_1(t), \dots, \pi_M(t)]$ be a $1 \times M$ row vector such that $\pi_i(t)$ equal to the $P[X(t) = s_i]$. Therefore, we have:

$$\pi(t) = \pi(0)\mathbf{\Pi}(t) \quad (5)$$

In general, finding $\pi(t)$ involves finding the corresponding eigenvalues and eigenvectors of \mathbf{Q} , which is *computationally difficult*.

Outline

- 1 Motivation
- 2 General Solution
- 3 Uniformization Method**
 - Uniformization
 - Probabilistic Interpretation
- 4 Practical Issues
 - Finite Sum
 - Extension to non-homogeneous CTMC

Uniformization is a computationally efficient method of performing transient analysis. Given a homogeneous CTMC $X(t)$ and the corresponding rate matrix \mathbf{Q} , let us define a homogeneous DTMC $X'(n)$ with the one-step transition probability matrix \mathbf{P} as:

$$\mathbf{P} = \mathbf{I} + \frac{\mathbf{Q}}{\Lambda} \quad (6)$$

where $\Lambda \geq \max_i \{q_i\}$, i.e., Λ is greater than or equal to the absolute diagonal value in \mathbf{Q} . Therefore, we have:

$$\mathbf{\Pi}(t) = \mathbf{e}^{\mathbf{Q}t} = \mathbf{e}^{(\mathbf{P}-\mathbf{I})\Lambda t} = \mathbf{e}^{\mathbf{P}\Lambda t} \mathbf{e}^{-\Lambda t} = \sum_{n=0}^{\infty} \mathbf{P}^n \frac{(\Lambda t)^n}{n!} \mathbf{e}^{-\Lambda t} \quad (7)$$

$$\pi(t) = \pi(0)\mathbf{\Pi}(t) = \sum_{n=0}^{\infty} \pi(0) \mathbf{P}^n \frac{(\Lambda t)^n}{n!} \mathbf{e}^{-\Lambda t} = \sum_{n=0}^{\infty} \pi(n) \frac{(\Lambda t)^n}{n!} \mathbf{e}^{-\Lambda t} \quad (8)$$

Consider a homogeneous CTMC X with state space S with M states and with rate matrix \mathbf{Q} . The diagonal element q_i in \mathbf{Q} represents the *output* rate (which is exponential). That is, once in s_i , the process moves to a different state with an exponential rate q_i . Furthermore, when it moves to other state, it moves to state s_j with probability q_{ij}/q_i .

DTMC Construction

We construct a DTMC X' with one step transition probability matrix \mathbf{P} (where $\mathbf{P} = \mathbf{I} + \mathbf{Q}/\Lambda$). We construct X' from X such that:

- X' has the same state space as X .
- The residence time in *any* state before a transition occurs is exponential with rate Λ .
- The probability that X' make a transition from s_i to s_j ($i \neq j$) is equal to q_{ij}/Λ . Furthermore, X' make a transition back to the same state with probability $(1 - q_i/\Lambda)$.

Consider a homogeneous CTMC X with state space S with M states and with rate matrix \mathbf{Q} . The diagonal element q_i in \mathbf{Q} represents the *output* rate (which is exponential). That is, once in s_i , the process moves to a different state with an exponential rate q_i . Furthermore, when it moves to other state, it moves to state s_j with probability q_{ij}/q_i .

DTMC Construction

We construct a DTMC X' with one step transition probability matrix \mathbf{P} (where $\mathbf{P} = \mathbf{I} + \mathbf{Q}/\Lambda$). We construct X' from X such that:

- X' has the same state space as X .
- The residence time in *any* state before a transition occurs is exponential with rate Λ .
- The probability that X' make a transition from s_i to s_j ($i \neq j$) is equal to q_{ij}/Λ . Furthermore, X' make a transition back to the same state with probability $(1 - q_i/\Lambda)$.

Consider a homogeneous CTMC X with state space S with M states and with rate matrix \mathbf{Q} . The diagonal element q_i in \mathbf{Q} represents the *output* rate (which is exponential). That is, once in s_i , the process moves to a different state with an exponential rate q_i . Furthermore, when it moves to other state, it moves to state s_j with probability q_{ij}/q_i .

DTMC Construction

We construct a DTMC X' with one step transition probability matrix \mathbf{P} (where $\mathbf{P} = \mathbf{I} + \mathbf{Q}/\Lambda$). We construct X' from X such that:

- X' has the same state space as X .
- The residence time in *any* state before a transition occurs is exponential with rate Λ .
- The probability that X' make a transition from s_i to s_j ($i \neq j$) is equal to q_{ij}/Λ . Furthermore, X' make a transition back to the same state with probability $(1 - q_i/\Lambda)$.

Consider a homogeneous CTMC X with state space S with M states and with rate matrix \mathbf{Q} . The diagonal element q_i in \mathbf{Q} represents the *output* rate (which is exponential). That is, once in s_i , the process moves to a different state with an exponential rate q_i . Furthermore, when it moves to other state, it moves to state s_j with probability q_{ij}/q_i .

DTMC Construction

We construct a DTMC X' with one step transition probability matrix \mathbf{P} (where $\mathbf{P} = \mathbf{I} + \mathbf{Q}/\Lambda$). We construct X' from X such that:

- X' has the same state space as X .
- The residence time in *any* state before a transition occurs is exponential with rate Λ .
- The probability that X' make a transition from s_i to s_j ($i \neq j$) is equal to q_{ij}/Λ . Furthermore, X' make a transition back to the same state with probability $(1 - q_i/\Lambda)$.

First Observation

Note that the probability that X' moves from s_i to s_j (for $i \neq j$) given that the transition is for a different state from s_i is:

$$\frac{q_{ij}/\Lambda}{q_i/\Lambda} = \frac{q_{ij}}{q_i},$$

which is equal to the probability that X goes from s_i to s_j in a transition.

Second Observation

- Given that the process makes n transitions to the same state s_i before leaving it, the distribution of the residence time in this state is simply the sum of n random variables with exponential distribution and rate Λ . This is equal to the Erlangian- n distribution and the density function $E'_{n,\Lambda}(t)$ is given by:

$$E'_{n,\Lambda}(t) = \frac{\Lambda (\Lambda t)^{n-1} e^{-\Lambda t}}{(n-1)!}$$

- The probability that the process makes $n-1$ self transitions before leaving s_i is:

$$\left(1 - \frac{q_i}{\Lambda}\right)^{n-1} \frac{q_i}{\Lambda}$$

- Combining the last two expressions (via theorem of total probability), then we see that the *total residence time* in s_i has an exponential distribution with rate q_i .

Second Observation

- Given that the process makes n transitions to the same state s_i before leaving it, the distribution of the residence time in this state is simply the sum of n random variables with exponential distribution and rate Λ . This is equal to the Erlangian- n distribution and the density function $E'_{n,\Lambda}(t)$ is given by:

$$E'_{n,\Lambda}(t) = \frac{\Lambda (\Lambda t)^{n-1} e^{-\Lambda t}}{(n-1)!}$$

- The probability that the process makes $n - 1$ self transitions before leaving s_i is:

$$\left(1 - \frac{q_i}{\Lambda}\right)^{n-1} \frac{q_i}{\Lambda}$$

- Combining the last two expressions (via theorem of total probability), then we see that the *total residence time* in s_i has an exponential distribution with rate q_i .

Second Observation

- Given that the process makes n transitions to the same state s_i before leaving it, the distribution of the residence time in this state is simply the sum of n random variables with exponential distribution and rate Λ . This is equal to the Erlangian- n distribution and the density function $E'_{n,\Lambda}(t)$ is given by:

$$E'_{n,\Lambda}(t) = \frac{\Lambda (\Lambda t)^{n-1} e^{-\Lambda t}}{(n-1)!}$$

- The probability that the process makes $n - 1$ self transitions before leaving s_i is:

$$\left(1 - \frac{q_i}{\Lambda}\right)^{n-1} \frac{q_i}{\Lambda}$$

- Combining the last two expression (via theorem of total probability), then we see that the *total residence time* in s_i has an exponential distribution with rate q_i .

Third Observation

Since the total residence time and transition probabilities of X and X' are the same, we can make the following conclusion:

$$\pi'_{ij}(t) = P[X'(t) = s_j | X'(0) = s_i] = P[X(t) = s_j | X(0) = s_i] = \pi_{ij}(t)$$

Therefore, the process CTMC X is *equivalent* to the DTMC X' subordinated to a Poisson process.

Outline

- 1 Motivation
- 2 General Solution
- 3 Uniformization Method
 - Uniformization
 - Probabilistic Interpretation
- 4 Practical Issues**
 - Finite Sum
 - Extension to non-homogeneous CTMC

Computational Procedure

Therefore, given \mathbf{Q} , we need to find $\mathbf{\Pi}(t)$. We first get \mathbf{P} , then solve for $\pi(n) = \pi(0)\mathbf{P}^n$ and then *weight* it by the probability that in the interval t , there are n Poisson arrival events.

$$\pi(t) = \pi(0)\mathbf{\Pi}(t) = \sum_{n=0}^{\infty} \pi(0)\mathbf{P}^n \frac{(\Lambda t)^n}{n!} e^{-\Lambda t} = \sum_{n=0}^{\infty} \pi(n) \frac{(\Lambda t)^n}{n!} e^{-\Lambda t} \quad (9)$$

So, given \mathbf{P} , find $\pi(n)$, which is just a vector matrix multiplication. Since in most cases, \mathbf{P} is a *sparse matrix*, therefore, it is efficient to compute. Then weighted the X' via an Poisson probability that there will be n Poisson arrival events.

To avoid the infinite summing series, we can use *truncation* and at the same time, know the error in advance! For example, if we only consider $N + 1$ transitions, we have:

$$\pi(t) = \sum_{n=0}^N \pi(n) e^{-\Lambda t} \frac{(\Lambda t)^n}{n!} + \epsilon(N) \quad (10)$$

where $\epsilon(N)$ is the error when the series is truncated after N terms. One important advantage of the uniformization method is that it is possible to find N in advance for a given tolerance since $\|\pi(n)\|_{\infty} \leq 1$ or:

$$\begin{aligned} \epsilon(N) &= \sum_{n=N+1}^{\infty} \pi(n) e^{-\Lambda t} \frac{(\Lambda t)^n}{n!} \\ &\leq \sum_{n=N+1}^{\infty} e^{-\Lambda t} \frac{(\Lambda t)^n}{n!} = 1 - \sum_{n=0}^N e^{-\Lambda t} \frac{(\Lambda t)^n}{n!} \end{aligned} \quad (11)$$

Non-homogeneous CTMC

Consider a non-homogeneous CTMC that has *different* transition rates for different interval of times. That is, for each interval $[t_{i-1}, t_i)$, the corresponding rate matrix is \mathbf{Q}_i (we are assuming that X_i to be time homogeneous). To extend this transient solution technique, let \mathbf{P}_i be the transition probability matrix of process X_i after uniformization, we have:

$$\pi(t) = \sum_{n=0}^{\infty} \pi_i(n) e^{-\Lambda_i t} \frac{(\Lambda_i t)^n}{n!} \quad \text{for } t_{i-1} \leq t < t_i \quad (12)$$

$$\pi_i(n) = \pi_i(n-1) \mathbf{P}_i \quad (13)$$

$$\pi_i(0) = \pi(t_{i-1}) \quad (14)$$

Let ϵ be the component-wise error bound, we spread the error into all intervals (weighted) by:

$$\epsilon_j = \frac{t_j - t_{j-1}}{t} \epsilon \quad (15)$$

And N_j can be computed based on ϵ_j by:

$$\epsilon_j \leq 1 - \sum_{n=0}^{N_j} e^{-\Lambda_j t} \frac{(\Lambda_j t)^n}{n!} \quad (16)$$