

COMP3506/7505: Regular Exercise Set 2

Prepared by Yufei Tao

Problem 1. Prove $30\sqrt{n} = O(\sqrt{n})$.

Solution. Set $c_1 = 30$ and $c_2 = 1$. The inequality $30\sqrt{n} \leq c_1 n$ holds for all $n \geq c_2$. This completes the proof.

Problem 2. Prove $\sqrt{n} = O(n)$.

Solution. Set $c_1 = 1$ and $c_2 = 1$. The inequality $\sqrt{n} \leq c_1 n$ holds for all $n \geq c_2$. This completes the proof.

Problem 3. Let $f(n)$, $g(n)$, and $h(n)$ be functions of integer n . Prove: if $f(n) = O(g(n))$ and $g(n) = O(h(n))$, then $f(n) = O(h(n))$.

Solution. Since $f(n) = O(g(n))$, there exist constants c_1, c_2 such that for all $n \geq c_2$, it holds that

$$f(n) \leq c_1 g(n).$$

Similarly, since $g(n) = O(h(n))$, there exist constants c'_1, c'_2 such that for all $n \geq c'_2$, it holds that

$$g(n) \leq c'_1 h(n).$$

Set $c''_1 = c_1 c'_1$ and $c''_2 = \max\{c_2, c'_2\}$. From the above, we know that for all $n \geq c''_2$, it holds that

$$f(n) \leq c_1 g(n) \leq c_1 c'_1 h(n) = c''_1 h(n).$$

Therefore, $f(n) = O(h(n))$.

Problem 4. Prove $(2n + 2)^3 = O(n^3)$.

Solution. Set $c_1 = 4^3$ and $c_2 = 1$. The inequality $(2n + 2)^3 \leq c_1 n^3$ holds for all $n \geq c_2$. This completes the proof.

Problem 5. Prove or disprove: $4^n = O(2^n)$.

Solution. Consider the ratio $4^n/2^n$, which equals 2^n . The ratio clearly goes to ∞ when n tends to ∞ . Therefore, the statement is incorrect.

Problem 6. Prove or disprove: $\frac{1}{n} = O(1)$.

Solution. Set $c_1 = 1$ and $c_2 = 1$. The inequality $1/n \leq c_1 \cdot 1$ holds for all $n \geq c_2$. This completes the proof.

Problem 7. Prove that if $k \log_2 k = \Theta(n)$, then $k = \Theta(n/\log n)$.

Solution. Since $k \log_2 k = O(n)$, there exist constants c_1, c_2 such that $k \log_2 k \leq c_1 n$ for all $n \geq c_2$. On the other hand, $k \log_2 k = \Omega(n)$ indicates the existence of constants c'_1, c'_2 such that $k \log_2 k \geq c'_1 n$ for all $n \geq c'_2$. Therefore, for all $n \geq \max\{c_2, c'_2\}$, we have:

$$c'_1 n \leq k \log_2 k \leq c_1 n. \quad (1)$$

Set $c''_2 = \max\{c_1, c_2, c'_2\}$.

When $n \geq c''_2$, we derive from (1):

$$\begin{aligned} \log_2(c'_1 n) &\leq \log_2(k \log_2 k) \leq \log_2(c_1 n) \\ \Rightarrow \log_2 c'_1 + \log_2 n &\leq \log_2 k + \log_2 \log_2 k \leq \log_2 c_1 + \log_2 n \\ \Rightarrow \begin{cases} \log_2 k \leq \log_2 c_1 + \log_2 n \leq 2 \log_2 n & (\text{using } n \geq c_1) \\ 2 \log_2 k \geq \log_2 k + \log_2 \log_2 k \geq \log_2 c'_1 + \log_2 n \geq \log_2 n \end{cases} \\ \Rightarrow \frac{\log_2 n}{2} &\leq \log_2 k \leq \log_2 n. \end{aligned} \quad (2)$$

Combining (1) and (2) leads to

$$\begin{cases} k \leq c_1 \frac{n}{\log_2 k} \leq 2c_1 \frac{n}{\log_2 n} \\ k \geq c'_1 \frac{n}{\log_2 k} \geq c'_1 \frac{n}{\log_2 n} \end{cases}$$

which means $k = \Theta(n / \log n)$.